arXiv:math/0006027v1 [math.AG] 5 Jun 2000

LOCAL COHOMOLOGY OF GENERALIZED OKAMOTO–PAINLEVÉ PAIRS AND PAINLEVÉ EQUATIONS

HITOMI TERAJIMA

Abstract. In the theory of deformation of Okamoto-Painlevé pair (S, Y), a local cohomology group $H^1_D(\Theta_S(-\log D))$ plays an important role. In this paper, we estimate the local cohomology group of pair (S, Y) for several types, and obtain the following results. For a pair (S, Y) corresponding to the space of initial conditions of the Painlevé equations, we show that the local cohomology group $H^1_D(\Theta_S(-\log D))$ is at least 1 dimensional. This fact is the key to understand Painlevé equation related to (S, Y). Moreover we show that, for the pairs (S, Y) of type $\tilde{A_8}$, the local cohomology group $H^1_D(\Theta_S(-\log D))$ vanish. Therefore in this case, there is no differential equation on $S - Y$ in the sense of the theory.

1. Introduction

To study the spaces of initial conditions of Painlevé equations constructed by Okamoto [O1] [O2] [O3], we introduced the notion of *generalized Okamoto-Painlevé pair* (S, Y) in [STT]. This is a pair of a complex projective surface S and an anti-canonical divisor $Y \in |-K_S|$ of S satisfying the following conditions: For the irreducible decomposition $Y = \sum_{i=1}^r m_i Y_i$, one has $Y \cdot Y_i = \deg Y|_{Y_i} = 0$ for $1 \leq i \leq r$. In addition, if S is a rational surface, (S, Y) is called a *generalized rational Okamoto-Painlevé pair.*

The generalized rational Okamoto-Painlevé pairs of non-fibered type are classified into three types; elliptic, multiplicative and additive. Moreover, the generalized rational Okamoto-Painlevé pairs of additive type such that $D := Y_{red} = \sum_{i=1}^r Y_i$ is a divisor with only normal crossings correspond to the spaces of initial conditions of the Painlevé equations (cf. [STT] [Sakai]) as follows:

type of (S, Y)	$\tilde{E_8}$	$\tilde{E_7}$	$\tilde{D_8}$	$\tilde{D_7}$	$\tilde{D_6}$	$\tilde{E_6}$	$\tilde{D_5}$	$\tilde{D_4}$
Painlevé equations	P_I	P_{II}	$P_{III}^{\tilde{D_8}}$	$P_{III}^{\tilde{D_7}}$	$P_{III} = P_{III}^{\tilde{D_6}}$	P_{IV}	P_V	P_{VI}

In what follows, (S, Y) is a generalized rational Okamoto-Painlevé pair of non-fibered type satisfying the condition: $D = Y_{red}$ is a normal crossing divisor with at least two irreducible components so that all irreducible components of Y_{red} are smooth rational curves. Let $\Theta_S(-\log D)$ be the sheaf of regular vector fields which have logarithmic zero along D. Here and after, all sheaves of $\mathcal{O}_S$-modules are considered in algebraic category.

1991 *Mathematics Subject Classification.* 14D15, 14J26, 32G10, 34M55.

Key words and phrases. Okamoto-Painlevé pair, local cohomology, Čech cohomology, the spaces of initial conditions of Painlevé equations.

Then we have the following key exact sequence:

$$0 \to H^1_D(\Theta_S(-\log D)) \to H^1(S, \Theta_S(-\log D)) \overset{res}{\to} H^1(S-D, \Theta_S(-\log D)),$$

where $H^1(S, \Theta_S(-\log D))$ and $H^1(S-D, \Theta_S(-\log D))$ are the space of infinitesimal deformations of the pair (S, Y) (cf. [Kaw]) and the space of infinitesimal deformations of $S-Y$, respectively. In [STT], we show that the directions corresponding to local cohomology $H^1_D(\Theta_S(-\log D))$ in the deformation of a pair (S, Y) induce differential equations on $S-Y$, by generalizing the Kodaira-Spencer theory to the open surface $S-Y$.

In this paper, we will show

$$\dim H^1_D(\Theta_S(-\log D)) \geq 1,$$

when (S, Y) is of additive type with the normal crossing divisor $D = Y_{red}$ (Theorem 2.1). This result is natural since $S-Y$ corresponds to the space of initial conditions of the Painlevé equations in this case. On the other hand, this is not always the case. In fact, we will prove that

$$H^1_D(\Theta_S(-\log D)) = \{0\},$$

for pairs (S, Y) of type $\tilde{A}_8$ which is classified as a multiplicative type (Proposition 4.1). This means that there does not exist differential equation on $S-Y$ by the way above. All these computations are carried out through calculations of Čech cohomologies by taking a coordinate system explicitly.

2. Local cohomology sequences and Time variables

Let (S, Y) be a generalized rational Okamoto–Painlevé pair and set $D = Y_{red}$. Moreover, in this section, we assume that

1. (S, Y) is of non-fibered type and
2. Y_{red} is a simple normal crossing divisor with at least two irreducible components, i.e. $(r \geq 2)$ so that all irreducible components of Y_{red} are smooth rational curves.

Here (S, Y) is called of fibered type if S has a structure of an elliptic surfacefibration $f : S \to \mathbf{P}^1$ with $f^*(\infty) = nY$ for some $n \geq 1$. If (S, Y) is not of fibered type, we call (S, Y) of non-fibered type. (cf. Definition 1.3, [STT]).

In what follows, $\mathcal{O}_S$ and $\mathcal{O}_{S-D}$ denote the sheaves of germs of algebraic regular functions on S and $S-D$ respectively. Moreover all sheaves of $\mathcal{O}_S$-modules are considered in algebraic category unless otherwise stated. Let us consider the following exact sequence of local cohomology groups ([Corollary 1.9, [Gr]])

$$\begin{array}{ccccccc} H^0(S, \Theta_S(-\log D)) & \to & H^0(S-D, \Theta_S(-\log D)) & \to & H^1_D(\Theta_S(-\log D)) & \to \\ H^1(S, \Theta_S(-\log D)) & \overset{\mu}{\to} & H^1(S-D, \Theta_S(-\log D)). & & & \end{array}$$

Since (S, Y) is of non-fibered type, from [(2), Proposition 2.1 [STT]], we see that

$$H^0(S-D, \Theta_S(-\log D)) = H^0(S-D, \Theta_S) = \{0\}.$$

Hence we have the following exact sequence:

$$0 \to \quad H^1_D(\Theta_S(-\log D)) \quad \to \quad H^1(S, \Theta_S(-\log D)) \quad \overset{\mu}{\to} \quad H^1(S-D, \Theta_S(-\log D)). \tag{1}$$

The following theorem is the main statement in this paper.

Theorem 2.1. *Let (S,Y) be a generalized rational Okamoto-Painlevé pair (S,Y) with the condition above. Moreover assume that $D=Y_{red}$ is of additive type. Then we have*

$$\dim H^0(D, \Theta_S(-\log D)\otimes N_D) = 1.$$

Here we put $N_D = \mathcal{O}_S(D)/\mathcal{O}_S$.

Since we have a natural inclusion

$$H^0(D, \Theta_S(-\log D)\otimes N_D) \hookrightarrow H^1_D(\Theta_S(-\log D)),$$

we obtain

$$\dim H^1_D(\Theta_S(-\log D)) \geq 1.$$

This theorem plays an important role to understand the Painlevé equation related to (S,Y).(cf. [STT]). Though we will not investigate the further structure of local cohomology here. Instead, we propose the following

Conjecture 2.1. *Under the same notation and assumption as in Theorem 2.1,*

$$H^1_D(\Theta_S(-\log D)) \simeq H^0(D, \Theta_S(-\log D)\otimes N_D) \simeq \mathbf{C}.$$

From the exact sequence (1), we see that the subspace $H^1_D(S,\Theta_S(-\log D))$ of $H^1(S,\Theta_S(-\log D))$ coincides with the kernel of μ. This implies that:

$$H^1_D(S,\Theta_S(-\log D)) \simeq \left\{ \begin{array}{c} \text{Infinitesimal deformations of } (S,D) \text{ whose restriction} \\ \text{to } S-D \text{ induces the trivial deformation} \end{array} \right\}.$$

In [§6 [STT]], we show that any direction corresponding to a non-zero element of the local cohomology group $H^1_D(S,\Theta_S(-\log D))$ induces a differential equation (at least locally) by using Čech coboundaries.

At this moment, we can not prove Conjecture 2.1 with full generality. However, we see that the one dimensional vector subspace $H^1(D,\Theta_S(-\log D)\otimes N_D)$ of $H^1_D(\Theta_S(-\log D)) \subset H^1(\Theta_S(-\log D))$ really corresponds to the time variable t in the known Painlevé equation. It is unlikely that we will have more time variables, so this gives an evidence of Conjecture 2.1.

Remark 2.1. We will consider (S,Y) of multiplicative type later, where the situation is different. (cf. Proposition 4.1).

Let us make preparations for the proof of Theorem 2.1.

Recall that

$$H^1_D(S,\Theta_S(-\log D)) = \varinjlim \operatorname{Ext}^1(\mathcal{O}_{nD}, \Theta_S(-\log D))$$

where $\mathcal{O}_{nD} = \mathcal{O}_S/\mathcal{O}_S(-nD)$ (cf. [Theorem 2.8, [Gr]]).

On the other hand, since $\Theta_S(-\log D)$ is a locally free $\mathcal{O}_S$-module, we see that

$$\mathcal{H}om(\mathcal{O}_{nD}, \Theta_S(-\log D)) = 0,$$

and

$$\mathcal{E}xt^1(\mathcal{O}_{nD}, \Theta_S(-\log D)) = \Theta_S(-\log D)\otimes N_{nD},$$

where $N_{nD} = \mathcal{O}_S(nD)/\mathcal{O}_S$. By an argument using a spectral sequence, we see that

$$H^1_D(S, \Theta_S(-\log D)) = \varinjlim H^0(\Theta_S(-\log D) \otimes N_{nD}).$$

Hence we have a natural inclusion

$$H^0(\Theta_S(-\log D) \otimes N_D) \hookrightarrow H^1_D(S, \Theta_S(-\log D)).$$

Lemma 2.1. *Let (S, Y) be a generalized rational Okamoto–Painlevé pair as above and set $D = Y_{red}$. Then we have the following exact sequences :*

$$0 \longrightarrow \Theta_D \otimes N_D \longrightarrow \Theta_S \otimes N_D \longrightarrow \nu_*(\oplus_{i=1}^r N_{Y_i/S}) \otimes N_D \longrightarrow 0, \tag{2}$$

$$0 \longrightarrow \nu_*(\oplus_{i=1}^r N_{Y_i/S}) \longrightarrow \Theta_S(-\log D) \otimes N_D \longrightarrow \Theta_D \otimes N_D \longrightarrow 0. \tag{3}$$

Here Θ_D denotes the tangent sheaf of D and $\nu : \tilde{D} = \coprod_{i=1}^9 Y_i \to D$ the normalization map.

Proof. The first exact sequence (2) follows from [(1.9), [B-W]].

Let us consider the following diagram:

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & \ker\lambda \arrow[d] & \\
0 \arrow[r] & \Theta_S(-\log D) \arrow[r] \arrow[d] & \Theta_S(-\log D) \otimes \mathcal{O}_S(D) \arrow[r] \arrow[d] & \Theta_S(-\log D) \otimes N_D \arrow[r] \arrow[d, "\lambda"] & 0 \\
0 \arrow[r] & \Theta_S \arrow[r] \arrow[d] & \Theta_S \otimes \mathcal{O}_S(D) \arrow[r] \arrow[d] & \Theta_S \otimes N_D \arrow[r] \arrow[d] & 0 \\
 & \nu_*(\oplus_{i=1}^r N_{Y_i/S}) \arrow[r, "\mu"] \arrow[d] & \nu_*(\oplus_{i=1}^r N_{Y_i/S}) \otimes N_D \arrow[r] \arrow[d] & \operatorname{coker}\lambda \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 & .
\end{tikzcd}$$

By the snake lemma, we obtain the exact sequence

$$0 \longrightarrow \ker\lambda \longrightarrow \nu_*(\oplus_{i=1}^r N_{Y_i/S}) \xrightarrow{\mu} \nu_*(\oplus_{i=1}^r N_{Y_i/S}) \otimes N_D \longrightarrow \operatorname{coker}\lambda \longrightarrow 0.$$

From a local consideration, we see that the map μ is a zero map, hence

$$\ker\lambda \simeq \nu_*(\oplus_{i=1}^r N_{Y_i/S}), \quad \text{and} \quad \nu_*(\oplus_{i=1}^r N_{Y_i/S}) \otimes N_D \simeq \operatorname{coker}\lambda.$$

Moreover since $\operatorname{Im}\lambda \simeq \ker[\Theta_S \otimes N_D \longrightarrow \nu_*(\oplus_{i=1}^r N_{Y_i/S}) \otimes N_D]$, from the exact sequence (2), we obtain the exact sequence (3). □

Note that since $N_{Y_i/S} = \mathcal{O}_{Y_i}(-2)$, we have

$$H^0(\oplus_{i=1}^r N_{Y_i/S}) = \{0\}, \quad H^1(\oplus_{i=1}^r N_{Y_i/S}) \simeq \mathbf{C}^r.$$

Moreover one can easily see that

$$\Theta_D \simeq \nu_*(\oplus_{i=1}^r \Theta_{Y_i}(-t_i)) \simeq \nu_*((\oplus_{i=1}^r \mathcal{O}_{Y_i}(2 - t_i))$$

where t_i is the number of intersections of Y_i with the other Y_j. On the other hand, since $D{\cdot}Y_i = t_i - 2$ and ν is a finite morphism, we see that

$$\begin{aligned} H^0(D, \Theta_D \otimes N_D) &\simeq H^0(D, \nu_*(\oplus_{i=1}^r \Theta_{Y_i}(-t_i)) \otimes N_D) \\ &\simeq H^0(\tilde{D}, (\oplus_{i=1}^r \Theta_{Y_i}(-t_i)) \otimes \nu^*(N_D)) \\ &\simeq \oplus_{i=1}^r H^0(Y_i, \Theta_{Y_i}(-t_i) \otimes N_D) \\ &\simeq \oplus_{i=1}^r H^0(Y_i, \mathcal{O}_{Y_i}) \\ &\simeq \mathbf{C}^r. \end{aligned}$$

Proof of Theorem 2.1.

From the exact sequence (3), one can obtain

$$H^0(\oplus_{i=1}^r N_{Y_i/S}) \longrightarrow H^0(\Theta_S(-\log D) \otimes N_D) \longrightarrow H^0(\Theta_D \otimes N_D) \xrightarrow{\delta} H^1(\oplus_{i=1}^r N_{Y_i/S})$$

where δ denotes the connected homomorphism.

From this sequence, the connecting homomorphism δ

$$\delta : H^0(\Theta_D \otimes N_D) \longrightarrow \oplus_{i=1}^r H^1(N_{Y_i/S})$$

can be identified with a linear map $\delta : \mathbf{C}^r \longrightarrow \mathbf{C}^r$ and we have an isomorphism

$$H^0(D, \Theta_S(-\log D) \otimes N_D) \simeq \ker \delta.$$

Now we state the following proposition, the proof of which is given in §3.

Proposition 2.1. *Let (S, Y) be a generalized rational Okamoto–Painlevé pair of additive type and set $D = Y_{red}$. Assume that D is a normal crossing divisor, then we can choose a basis of $H^0(\Theta_D \otimes N_D)$ and $\oplus_{i=1}^r H^1(N_{Y_i})$ so that the linear map $\delta : H^0(\Theta_D \otimes N_D) \to \oplus_{i=1}^r H^1(N_{Y_i})$ is represented by the intersection matrix $((Y_i \cdot Y_j))_{1\le i,j\le r}$ with respect to these basis.*

In this case, it is well-known that the rank of the intersection matrix $((Y_i \cdot Y_j))_{1\le i,j \le r}$ is $r-1$. Hence from the Proposition 2.1, we have

$$\dim_{\mathbf{C}} H^0(\Theta_S(-\log D) \otimes N_D) = \dim_{\mathbf{C}} \ker \delta = 1.$$

□

3. PROOF OF PROPOSITION 2.1

In this section, we shall prove Proposition 2.1.

Here we give a detailed proof only for the case of $\tilde{E}_7$. The proof of other cases are similar.

Let (S, Y) be a generalized rational Okamoto–Painlevé pair of type $\tilde{E}_7$. Then according to the results in Appendix B of [Sakai], (S, Y) can be obtained by blowings up 9-points of $\mathbf{P}^2$ as follows.

Let $[x:y:z]$ be the homogeneous coordinates of $\mathbf{P}^2$.


$z=0$
$x=0$
$p_1 \leftarrow p_2 \leftarrow p_3 \leftarrow p_6 \leftarrow$
$\leftarrow p_7 \leftarrow p_8 \leftarrow p_9$
p_5
p_4
$\longleftarrow$
p_9
p_5
p_4

$$
\begin{aligned}
&p_1 : (0:1:0) \leftarrow p_2 : \left(\frac{x}{y}, \frac{z}{x}\right) = (0,0) \leftarrow p_3 : \left(\frac{x}{y}, \frac{yz}{x^2}\right) = (0,0) \leftarrow p_6 : \left(\frac{x}{y}, \frac{y^2z}{x^3}\right) = (0,1) \leftarrow \\
&p_7 : \left(\frac{x}{y}, \frac{y(y^2z-x^3)}{x^4}\right) = (0,0) \leftarrow p_8 : \left(\frac{x}{y}, \frac{y^2(y^2z-x^3)}{x^5}\right) = (0,-s) \leftarrow \\
&p_9 : \left(\frac{x}{y}, \frac{y(y^2(y^2z-x^3)+sx^5)}{x^6}\right) = (0,-\alpha_0),
\end{aligned}
$$

$$
\begin{aligned}
&p_4 : (0:0:1), \\
&p_5 : (0:\alpha_1:1).
\end{aligned}
$$

Note that there exist three deformation parameters (α_0, α_1, s) of the blowings-up. Moreover there exists a $\mathbf{C}^{\times}$-action on the family of surfaces by

$$(\mu, (\alpha_0, \alpha_1; s, [x:y:z])) \mapsto (\mu^3\alpha_0, \mu^3\alpha_1, \mu^2 s; [x:\mu y:\mu^{-2}z]).$$

If we set $\lambda = \alpha_0 + \alpha_1$ and $\mu = s/\lambda$, then

$$t = \frac{s^3}{\lambda^2},\ a_0 = \frac{s^3\alpha_0}{\lambda^3},\ a_1 = \frac{s^3\alpha_1}{\lambda^3}, X = x,\ Y = \frac{sy}{\lambda},\ Z = \frac{\lambda^2 z}{s^2}$$

are invariant under the $\mathbf{C}^{\times}$-action, and we have the relation

$$a_0 + a_1 = t.$$

Now we can introduce the affine open covering of S and affine coordinates by the explicit blowings-up of $\mathbf{P}^2$. The following diagram shows how one can perform blowing-ups and introduce the new coordinates (x_i, y_i) $(1 \le i \le 13)$, (u_j, v_j) $(j = 14, 15, 16)$.



$$E_i \simeq \mathbf{P}^1,\ E_i^2 = -1,$$
$$Y_i \simeq \mathbf{P}^1,\ Y_i^2 = -2.$$

$$\begin{aligned}
U_i &= \operatorname{Spec}\mathbf{C}\left[x_i, y_i\right] \cong \mathbf{C}^2 \quad (i = 1, 2, \cdots, 7).\\
U_8 &= \operatorname{Spec}\mathbf{C}\left[x_8, y_8, \frac{1}{1+x_8}\right] \cong \mathbf{C}^2 - \{1 + x_8 = 0\}.\\
U_9 &= \operatorname{Spec}\mathbf{C}\left[x_9, y_9, \frac{1}{1+{x_9}^2 y_9}\right] \cong \mathbf{C}^2 - \{1 + {x_9}^2 y_9 = 0\}.\\
U_{10} &= \operatorname{Spec}\mathbf{C}\left[x_{10}, y_{10}, \frac{1}{1+x_{10}{y_{10}}^2}\right] \cong \mathbf{C}^2 - \{1 + x_{10}{y_{10}}^2 = 0\}.\\
U_{11} &= \operatorname{Spec}\mathbf{C}\left[x_{11}, y_{11}, \frac{1}{1-t{x_{11}}^2{y_{11}}^2 + {x_{11}}^3{y_{11}}^2}\right] \cong \mathbf{C}^2 - \{1 - t{x_{11}}^2{y_{11}}^2 + {x_{11}}^3{y_{11}}^2 = 0\}.\\
U_{12} &= \operatorname{Spec}\mathbf{C}\left[x_{12}, y_{12}, \frac{1}{1-t{y_{12}}^2 + x_{12}{y_{12}}^3}\right] \cong \mathbf{C}^2 - \{1 - t{y_{12}}^2 + x_{12}{y_{12}}^3 = 0\}.\\
U_{13} &= \operatorname{Spec}\mathbf{C}\left[x_{13}, y_{13}, \frac{1}{-1+t{x_{13}}^2{y_{13}}^2 + a_0{x_{13}}^3{y_{13}}^3 - {x_{13}}^4{y_{13}}^3}\right]\\
&\cong \mathbf{C}^2 - \{-1 + t{x_{13}}^2{y_{13}}^2 + a_0{x_{13}}^3{y_{13}}^3 - {x_{13}}^4{y_{13}}^3 = 0\}.\\
U_j &= \operatorname{Spec}\mathbf{C}\left[u_j, v_j\right] \cong \mathbf{C}^2 \quad (j = 14, 15).\\
U_{16} &= \operatorname{Spec}\mathbf{C}\left[u_{16}, v_{16}, \frac{1}{-1+tu_{16}^2 + a_0u_{16}^3 - u_{16}^4v_{16}}\right] \cong \mathbf{C}^2 - \{-1 + tu_{16}^2 + a_0u_{16}^3 - u_{16}^4v_3 = 0\}.
\end{aligned}$$

$$\begin{aligned}
&Y_1 = \{x_1 = 0, x_2 = 0, y_3 = 0\}, && Y_2 = \{x_3 = 0, y_4 = 0\},\\
&Y_3 = \{x_4 = 0, y_5 = 0\}, && Y_4 = \{x_5 = 0, y_6 = 0, y_8 = 0\},\\
&Y_5 = \{x_6 = 0, y_7 = 0\}, && Y_6 = \{x_8 = 0, y_9 = 0\},\\
&Y_7 = \{x_9 = 0, y_{10} = 0, y_{11} = 0\}, && Y_8 = \{x_{11} = 0, y_{12} = 0, y_{13} = 0\}.
\end{aligned}$$

$$S = \bigcup_{i=1}^{16} U_i.$$

$$Y = Y_1 + 2Y_2 + 3Y_3 + 4Y_4 + 2Y_5 + 3Y_6 + 2Y_7 + Y_8, \quad D = \sum_{i=1}^{8} Y_i.$$

$$S - Y = U_{14} \cup U_{15} \cup U_{16}.$$

More explicitly, new coordinates can be given by the following formula

$$\begin{cases} x_1 = \dfrac{X}{Y} \\ y_1 = \dfrac{Y}{Z} \end{cases} \quad \begin{cases} x_2 = \dfrac{X}{Y - a_1 Z} \\ y_2 = \dfrac{Y - a_1 Z}{Z} \end{cases} \quad \begin{cases} x_3 = \dfrac{Z}{Y} \\ y_3 = \dfrac{X}{Z} \end{cases}$$

$$\begin{cases} x_4 = \dfrac{Z}{X} \\ y_4 = \dfrac{X^2}{YZ} \end{cases} \quad \begin{cases} x_5 = \dfrac{YZ}{X^2} \\ y_5 = \dfrac{X^3}{Y^2 Z} \end{cases} \quad \begin{cases} x_6 = \dfrac{Y^2 Z}{X^3} \\ y_6 = \dfrac{X}{Y} \end{cases}$$

$$\begin{cases} x_7 = \dfrac{Y}{X} \\ y_7 = \dfrac{Z}{X} \end{cases} \quad \begin{cases} x_8 = \dfrac{Y^2 Z - X^3}{X^3} \\ y_8 = \dfrac{X^4}{Y(Y^2 Z - X^3)} \end{cases} \quad \begin{cases} x_9 = \dfrac{Y(Y^2 Z - X^3)}{X^4} \\ y_9 = \dfrac{X^5}{Y^2(Y^2 Z - X^3)} \end{cases}$$

$$\begin{cases} x_{10} = \dfrac{Y^2(Y^2 Z - X^3)}{X^5} \\ y_{10} = \dfrac{X}{Y} \end{cases} \quad \begin{cases} x_{11} = \dfrac{Y^2(Y^2 Z - X^3) + tX^5}{X^5} \\ y_{11} = \dfrac{X^6}{Y(Y^2(Y^2 Z - X^3) + tX^5)} \end{cases}$$

$$\begin{cases} x_{12} = \dfrac{Y(Y^2(Y^2 Z - X^3) + tX^5)}{X^6} \\ y_{12} = \dfrac{X}{Y} \end{cases} \quad \begin{cases} x_{13} = \dfrac{Y(Y^2(Y^2 Z - X^3) + tX^5) + a_0 X^6}{X^6} \\ y_{13} = \dfrac{X^7}{Y(Y(Y^2(Y^2 Z - X^3) + tX^5) + a_0 X^6)} \end{cases}$$

$$\begin{cases} u_{14} = \dfrac{X}{Z} \\ v_{14} = \dfrac{Y}{X} \end{cases} \quad \begin{cases} u_{15} = \dfrac{X}{Z} \\ v_{15} = \dfrac{Y - a_1 Z}{X} \end{cases} \quad \begin{cases} u_{16} = \dfrac{X}{Y} \\ v_{16} = \dfrac{Y(Y(Y^2(Y^2 Z - X^3) + tX^5) + a_0 X^6)}{X^7}. \end{cases}$$

From these formula, we can determine the coordinate transformation between (x_i, y_i)'s and (u_j, v_j)'s. For later use, we need only the coordinate transformations near each component Y_i. Here we will list up the coordinate transformations only for a neighborhood of each Y_i.

$$Y_1 : \begin{cases} x_1 = \dfrac{x_2 y_2}{a_1 + y_2} \\ y_1 = a_1 + y_2 \end{cases} \begin{cases} x_1 = x_3 y_3 \\ y_1 = \dfrac{1}{x_3} \end{cases} \qquad Y_5 : \begin{cases} x_6 = {x_7}^2 y_7 \\ y_6 = \dfrac{1}{x_7} \end{cases}$$

$$Y_2 : \begin{cases} x_3 = {x_4}^2 y_4 \\ y_3 = \dfrac{1}{x_4} \end{cases} \qquad Y_6 : \begin{cases} x_8 = {x_9}^2 y_9 \\ y_8 = \dfrac{1}{x_9} \end{cases}$$

$$Y_3 : \begin{cases} x_4 = {x_5}^2 y_5 \\ y_4 = \dfrac{1}{x_5} \end{cases} \qquad Y_7 : \begin{cases} x_9 = x_{10} y_{10} \\ y_9 = \dfrac{1}{x_{10}} \end{cases} \begin{cases} x_9 = x_{11}(-t + x_{11}) y_{11} \\ y_9 = \dfrac{1}{-t + x_{11}} \end{cases}$$

$$Y_4 : \begin{cases} x_5 = x_6 y_6 \\ y_5 = \dfrac{1}{x_6} \end{cases} \begin{cases} x_5 = x_8(1 + x_8) y_8 \\ y_5 = \dfrac{1}{1 + x_8} \end{cases} \qquad Y_8 : \begin{cases} x_{11} = x_{12} y_{12} \\ y_{11} = \dfrac{1}{x_{12}} \end{cases} \begin{cases} x_{11} = x_{13}(-a_0 + x_{13}) y_{13} \\ y_{11} = \dfrac{1}{-a_0 + x_{13}}. \end{cases}$$

Let us consider the sheaf $\Theta_S(-\log D)$ and the sheaf exact sequence

$$0 \longrightarrow \nu_*(\oplus_{i=1}^{8} N_{Y_i/S}) \longrightarrow \Theta_S(-\log D) \otimes N_D \longrightarrow \Theta_D \otimes N_D \longrightarrow 0.$$

We will analyse the edge homomorphism

$$\delta : H^0(\Theta_D \otimes N_D) \longrightarrow H^1(\nu_*(\oplus_{i=1}^{8} N_{Y_i/S})) \tag{4}$$

by using the Čech cocycles.

Noting that ν is a finite morphism, and $\Theta_D \otimes N_D \simeq \nu_*(\oplus_{i=1}^{8} \Theta_{Y_i}(-t_i)) \otimes N_D$ where t_i is the number of intersections of Y_i with other components, we see that

$$H^0(D, \Theta_D \otimes N_D) \simeq H^0(D, \nu_*(\oplus_{i=1}^{8} \Theta_{Y_i}(-t_i)) \otimes N_D) \simeq \mathbf{C}^8. \tag{5}$$

For each i $(1 \leq i \leq 8)$, we introduce a generator θ_i of the cohomology group in (5) as follows.

θ_1	$\left\{\begin{array}{l} \theta_1^1 = \frac{-a_1+y_1}{x_1}\frac{\partial}{\partial y_1} \text{ on } U_1 \cap Y_1, \quad \theta_1^2 = \frac{a_1+y_2}{x_2}\frac{\partial}{\partial y_2} \text{ on } U_2 \cap Y_1, \\ \theta_1^3 = \frac{-1+a_1 x_3}{y_3}\frac{\partial}{\partial x_3} \text{ on } U_3 \cap Y_1 \end{array}\right\}$
θ_2	$\left\{\theta_2^3 = \frac{1}{x_3}\frac{\partial}{\partial y_3} \text{ on } U_3 \cap Y_2, \quad \theta_2^4 = -\frac{1}{y_4}\frac{\partial}{\partial x_4} \text{ on } U_4 \cap Y_2\right\}$
θ_3	$\left\{\theta_3^4 = \frac{1}{x_4}\frac{\partial}{\partial y_4} \text{ on } U_4 \cap Y_3, \quad \theta_3^5 = -\frac{1}{y_5}\frac{\partial}{\partial x_5} \text{ on } U_5 \cap Y_3\right\}$
θ_4	$\left\{\begin{array}{l} \theta_4^5 = \frac{1-y_5}{x_5}\frac{\partial}{\partial y_5} \text{ on } U_5 \cap Y_4, \quad \theta_4^6 = \frac{1-x_6}{y_6}\frac{\partial}{\partial x_6} \text{ on } U_6 \cap Y_4, \\ \theta_4^8 = -\frac{1}{y_8}\frac{\partial}{\partial x_8} \text{ on } U_8 \cap Y_4 \end{array}\right\}$
θ_5	$\left\{\theta_5^6 = -\frac{1}{x_6}\frac{\partial}{\partial y_6} \text{ on } U_6 \cap Y_5, \quad \theta_5^7 = \frac{1}{y_7}\frac{\partial}{\partial x_7} \text{ on } U_7 \cap Y_5\right\}$
θ_6	$\left\{\theta_6^8 = \frac{1}{x_8}\frac{\partial}{\partial y_8} \text{ on } U_8 \cap Y_6, \quad \theta_6^9 = -\frac{1}{y_9}\frac{\partial}{\partial x_9} \text{ on } U_8 \cap Y_6\right\}$
θ_7	$\left\{\begin{array}{l} \theta_7^9 = \frac{1+ty_9}{x_9}\frac{\partial}{\partial y_9} \text{ on } U_9 \cap Y_7, \quad \theta_7^{10} = -\frac{t+x_{10}}{y_{10}}\frac{\partial}{\partial x_{10}} \text{ on } U_{10} \cap Y_7, \\ \theta_7^{11} = -\frac{1}{y_{11}}\frac{\partial}{\partial x_{11}} \text{ on } U_{11} \cap Y_7 \end{array}\right\}$
θ_8	$\left\{\begin{array}{l} \theta_8^{11} = \frac{1+a_0 y_{11}}{x_{11}}\frac{\partial}{\partial y_{11}} \text{ on } U_{11} \cap Y_8, \quad \theta_8^{12} = -\frac{a_0+x_{12}}{y_{12}}\frac{\partial}{\partial x_{12}} \text{ on } U_{12} \cap Y_8, \\ \theta_8^{13} = -\frac{1}{y_{13}}\frac{\partial}{\partial x_{13}} \text{ on } U_{13} \cap Y_8 \end{array}\right\}$

On the other hand, for each i $(1 \leq i \leq 8)$, we have a generator $\eta_i \in H^1(Y_i, N_{Y_i/S})$ as follows.

η_1	$\left\{\eta_1^{12} = 0 \text{ on } U_1 \cap U_2 \cap Y_1, \quad \eta_1^{13} = \frac{1}{x_3}\frac{\partial}{\partial y_3} \text{ on } U_1 \cap U_3 \cap Y_1\right\}$
η_2	$\left\{\eta_2^{34} = \frac{1}{x_4}\frac{\partial}{\partial y_4} \text{ on } U_3 \cap U_4 \cap Y_2\right\}$
η_3	$\left\{\eta_3^{45} = \frac{1}{x_5}\frac{\partial}{\partial y_5} \text{ on } U_4 \cap U_5 \cap Y_3\right\}$
η_4	$\left\{\eta_4^{56} = 0 \text{ on } U_5 \cap U_6 \cap Y_4, \quad \eta_4^{58} = \frac{1}{x_8}\frac{\partial}{\partial y_8} \text{ on } U_5 \cap U_8 \cap Y_4\right\}$
η_5	$\left\{\eta_5^{67} = -\frac{1}{x_7}\frac{\partial}{\partial y_7} \text{ on } U_6 \cap U_7 \cap Y_5\right\}$
η_6	$\left\{\eta_6^{89} = \frac{1}{x_9}\frac{\partial}{\partial y_9} \text{ on } U_8 \cap U_9 \cap Y_6\right\}$
η_7	$\left\{\eta_7^{9\,10} = 0 \text{ on } U_9 \cap U_{10} \cap Y_7, \quad \eta_7^{9\,11} = \frac{1}{x_{11}}\frac{\partial}{\partial y_{11}} \text{ on } U_9 \cap U_{11} \cap Y_7\right\}$
η_8	$\left\{\eta_8^{11\,13} = \frac{1}{x_{13}}\frac{\partial}{\partial y_{13}} \text{ on } U_{11} \cap U_{13} \cap Y_8, \quad \eta_8^{12\,13} = 0 \text{ on } U_{12} \cap U_{13} \cap Y_8\right\}$

We take $\{\theta_i\}$ and $\{\eta_i\}$ as basis of $H^0(\Theta_D \otimes N_D)$ and $\oplus_{i=1}^{8} H^1(N_{Y_i/M})$ respectively.

By using these bases, we compute the matrix representing the connecting homomorphism δ. For that purpose, let us lift 0-cocycle θ_1 to 0-cochains of $\Theta_S(-\log D) \otimes N_D$ as

$$\tilde{\theta}_1^1 = \frac{-a_1 + y_1}{x_1}\frac{\partial}{\partial y_1} \text{ on } U_1, \quad \tilde{\theta}_1^2 = \frac{a_1 + y_2}{x_2}\frac{\partial}{\partial y_2} \text{ on } U_2, \quad \tilde{\theta}_1^3 = \frac{-1 + a_1 x_3}{y_3}\frac{\partial}{\partial x_3} \text{ on } U_3,$$
$$\tilde{\theta}_1^i = 0 \text{ on } U_i \ (i = 4, 5, \cdots, 16).$$

Other 0-cocycles can be lifted in a similar way.

We first compute $\delta(\theta_1)$.

From the definition of δ, we have $\delta(\theta_1) = \{\delta(\theta_1)_{ij} \text{ on } U_i \cap U_j \cap D\}$ with

$$\begin{cases} \delta(\theta_1)_{12} = (\tilde{\theta}_1^2 - \tilde{\theta}_1^1)|_{Y_1} = \left(\dfrac{a_1 + y_2}{x_2}\dfrac{\partial}{\partial y_2} - \dfrac{-a_1 + y_1}{x_1}\dfrac{\partial}{\partial y_1}\right)\Bigg|_{Y_1} = \dfrac{a_1}{y_2}\dfrac{\partial}{\partial x_2} \\ \delta(\theta_1)_{13} = (\tilde{\theta}_1^3 - \tilde{\theta}_1^1)|_{Y_1} = \left(\dfrac{-1 + a_1 x_3}{y_3}\dfrac{\partial}{\partial x_3} - \dfrac{-a_1 + y_1}{x_1}\dfrac{\partial}{\partial y_1}\right)\Bigg|_{Y_1} = \dfrac{-1 + a_1 x_3}{x_3}\dfrac{\partial}{\partial y_3} \end{cases}$$
$$\delta(\theta_1)_{34} = (\tilde{\theta}_1^4 - \tilde{\theta}_1^3)|_{Y_2} = \left(0 - \frac{-1 + a_1 x_3}{y_3}\frac{\partial}{\partial x_3}\right)\Bigg|_{Y_2} = \frac{1}{x_4}\frac{\partial}{\partial y_4} = \eta_2^{34}$$

Other $\delta(\theta_1)_{ij}$'s are zero.

Obviously $\{\delta(\theta_1)_{34}\} = \{\eta_2^{34}\} = \eta_2$.

In order to see that $\{\delta(\theta_1)_{12}, \delta(\theta_1)_{13}\} = -2\eta_1$, we set $\tau = \{\tau_1 = -\frac{\partial}{\partial x_1}, \tau_2 = -\frac{\partial}{\partial x_2}, \tau_3 = a_1 \frac{\partial}{\partial y_3}\} \in C^0(N_{Y_1/S})$.

$$\begin{cases} \delta(\theta_1)_{12} + 2\eta_1^{12} = \dfrac{a_1}{y_2}\dfrac{\partial}{\partial x_2} = \tau_2 - \tau_1 \\ \delta(\theta_1)_{13} + 2\eta_1^{13} = \dfrac{1 + a_1 x_3}{x_3}\dfrac{\partial}{\partial y_3} = \tau_3 - \tau_1 \end{cases}$$

This implies $\{\delta(\theta_1)_{12}, \delta(\theta_1)_{13}\} + 2\eta_1 = \delta\tau$. Then we have $\delta(\theta_1) = -2\eta_1 + \eta_2$.

Other $\delta(\theta_i)$'s can be treated in a similar way. In what follows, we just list up the results of computations.

○ $\delta(\theta_2) = \eta_1 - 2\eta_2 + \eta_3$

$$\begin{cases} \delta(\theta_2)_{12} = \tilde{\theta}_2^2 - \tilde{\theta}_2^1|_{Y_1} = 0 - 0 = \eta_1^{12} \\ \delta(\theta_2)_{13} = \tilde{\theta}_2^3 - \tilde{\theta}_2^1|_{Y_1} = \dfrac{1}{x_3}\dfrac{\partial}{\partial y_3} - 0 = \eta_1^{13} \end{cases}$$
$$\delta(\theta_2)_{34} = \tilde{\theta}_2^4 - \tilde{\theta}_2^3|_{Y_2} = -\frac{1}{y_4}\frac{\partial}{\partial x_4} - \frac{1}{x_3}\frac{\partial}{\partial y_3} = -2\frac{1}{x_4}\frac{\partial}{\partial y_4} = -2\eta_2^{34}$$
$$\delta(\theta_2)_{45} = \tilde{\theta}_2^5 - \tilde{\theta}_2^4|_{Y_3} = 0 - \left(-\frac{1}{y_4}\frac{\partial}{\partial x_4}\right) = \frac{1}{x_5}\frac{\partial}{\partial y_5} = \eta_3^{45}$$

$\circ\ \delta(\theta_3) = \eta_2 - 2\eta_3 + \eta_4$

$$\begin{aligned}
&\delta(\theta_3)_{34} = \tilde{\theta}_3^4 - \tilde{\theta}_3^3|_{Y_2} = \frac{1}{x_4}\frac{\partial}{\partial y_4} - 0 = \frac{1}{x_4}\frac{\partial}{\partial y_4} = \eta_2^{34} \\
&\delta(\theta_3)_{45} = \tilde{\theta}_3^5 - \tilde{\theta}_3^4|_{Y_3} = -\frac{1}{y_5}\frac{\partial}{\partial x_5} - \frac{1}{x_4}\frac{\partial}{\partial y_4} = -2\frac{1}{x_5}\frac{\partial}{\partial y_5} = -2\eta_3^{45} \\
&\begin{cases}
\delta(\theta_3)_{56} = \tilde{\theta}_3^6 - \tilde{\theta}_3^5|_{Y_4} = 0 - \left(-\dfrac{1}{y_5}\dfrac{\partial}{\partial x_5}\right) = \dfrac{\partial}{\partial y_6} \\
\delta(\theta_3)_{58} = \tilde{\theta}_3^8 - \tilde{\theta}_3^5|_{Y_4} = 0 - \left(-\dfrac{1}{y_5}\dfrac{\partial}{\partial x_5}\right) = \dfrac{1}{x_8}\dfrac{\partial}{\partial y_8}
\end{cases}
\end{aligned}$$

Set $\{\tau_5 = 0, \tau_6 = \frac{\partial}{\partial y_6}, \tau_8 = 0\} \in C^0(N_{Y_4/S})$.

$$\begin{cases}
\delta(\theta_3)_{56} - \eta_4^{56} = \dfrac{\partial}{\partial y_6} - 0 = \tau_6 - \tau_5 \\
\delta(\theta_3)_{58} - \eta_4^{58} = \dfrac{1}{x_8}\dfrac{\partial}{\partial y_8} - \dfrac{1}{x_8}\dfrac{\partial}{\partial y_8} = 0 = \tau_8 - \tau_5
\end{cases}$$

Thus we have $\{\delta(\theta_3)_{56}, \delta(\theta_3)_{58}\} = \eta_4$.

$\circ\ \delta(\theta_4) = \eta_3 - 2\eta_4 + \eta_5 + \eta_6$

$$\begin{aligned}
&\delta(\theta_4)_{45} = \tilde{\theta}_4^5 - \tilde{\theta}_4^4|_{Y_3} = \frac{1-y_5}{x_5}\frac{\partial}{\partial y_5} - 0 = \frac{1}{x_5}\frac{\partial}{\partial y_5} = \eta_3^{45} \\
&\begin{cases}
\delta(\theta_4)_{56} = \tilde{\theta}_4^6 - \tilde{\theta}_4^5|_{Y_4} = \dfrac{1-x_6}{y_6}\dfrac{\partial}{\partial x_6} - \dfrac{1-y_5}{x_5}\dfrac{\partial}{\partial y_5} = \dfrac{1-x_6}{x_6}\dfrac{\partial}{\partial y_6} \\
\delta(\theta_4)_{58} = \tilde{\theta}_4^8 - \tilde{\theta}_4^5|_{Y_4} = -\dfrac{1}{y_8}\dfrac{\partial}{\partial x_8} - \dfrac{1-y_5}{x_5}\dfrac{\partial}{\partial y_5} = -\dfrac{1+2x_8}{x_8(1+x_8)}\dfrac{\partial}{\partial y_8}
\end{cases} \\
&\delta(\theta_4)_{67} = \tilde{\theta}_4^7 - \tilde{\theta}_4^6|_{Y_5} = 0 - \frac{1-x_6}{y_6}\frac{\partial}{\partial x_6} = -\frac{1}{x_7}\frac{\partial}{\partial y_7} = \eta_5^{67} \\
&\delta(\theta_4)_{89} = \tilde{\theta}_4^9 - \tilde{\theta}_4^8|_{Y_6} = 0 - \left(-\frac{1}{y_8}\frac{\partial}{\partial x_8}\right) = \frac{1}{x_9}\frac{\partial}{\partial y_9} = \eta_6^{89}
\end{aligned}$$

Set $\{\tau_5 = -\frac{\partial}{\partial x_5}, \tau_6 = -\frac{\partial}{\partial y_6}, \tau_8 = 0\} \in C^0(N_{Y_4/S})$.

$$\begin{cases}
\delta(\theta_4)_{56} + 2\eta_4^{56} = \dfrac{1-x_6}{x_6}\dfrac{\partial}{\partial y_6} = \tau_6 - \tau_5 \\
\delta(\theta_4)_{58} + 2\eta_4^{56} = \dfrac{1}{x_8(1+x_8)}\dfrac{\partial}{\partial y_8} = \tau_8 - \tau_5
\end{cases}$$

Thus we have $\{\delta(\theta_4)_{56}, \delta(\theta_4)_{58}\} = -2\eta_4$.

∘ $\delta(\theta_5) = \eta_4 - 2\eta_5$

$$\begin{cases} \delta(\theta_5)_{56} = \tilde{\theta_5^6} - \tilde{\theta_5^5}|_{Y_4} = -\dfrac{1}{x_6}\dfrac{\partial}{\partial y_6} - 0 = -\dfrac{1}{x_6}\dfrac{\partial}{\partial y_6} \\ \delta(\theta_5)_{58} = \tilde{\theta_5^8} - \tilde{\theta_5^5}|_{Y_4} = 0 - 0 = 0 \end{cases}$$

$$\delta(\theta_5)_{67} = \tilde{\theta_5^7} - \tilde{\theta_5^6}|_{Y_5} = \frac{1}{y_7}\frac{\partial}{\partial x_7} - \left(-\frac{1}{x_6}\frac{\partial}{\partial y_6}\right) = 2\frac{1}{x_7}\frac{\partial}{\partial y_7} = -2\eta_5^{67}$$

Set $\{\tau_5 = \frac{\partial}{\partial x_5}, \tau_6 = 0, \tau_8 = 0\} \in C^0(N_{Y_4/S})$.

$$\begin{cases} \delta(\theta_5)_{56} - \eta_4^{56} = -\dfrac{1}{x_6}\dfrac{\partial}{\partial y_6} = \tau_6 - \tau_5 \\ \delta(\theta_5)_{58} - \eta_4^{58} = -\dfrac{1}{x_8(1+x_8)}\dfrac{\partial}{\partial y_8} = \tau_8 - \tau_5 \end{cases}$$

Thus we have $\{\delta(\theta_5)_{56}, \delta(\theta_5)_{58}\} = \eta_4$.

∘ $\delta(\theta_6) = \eta_4 - 2\eta_6 + \eta_7$

$$\begin{cases} \delta(\theta_6)_{56} = \tilde{\theta_6^6} - \tilde{\theta_6^5}|_{Y_4} = 0 - 0 = \eta_4^{56} \\ \delta(\theta_6)_{58} = \tilde{\theta_6^8} - \tilde{\theta_6^5}|_{Y_4} = \dfrac{1}{x_8}\dfrac{\partial}{\partial y_8} - 0 = \eta_4^{58} \end{cases}$$

$$\delta(\theta_6)_{89} = \tilde{\theta_6^9} - \tilde{\theta_6^8}|_{Y_6} = -\frac{1}{y_9}\frac{\partial}{\partial x_9} - \frac{1}{x_8}\frac{\partial}{\partial y_8} = -2\frac{1}{x_9}\frac{\partial}{\partial y_9} = -2\eta_6^{89}$$

$$\begin{cases} \delta(\theta_6)_{9\,10} = \tilde{\theta_6^{10}} - \tilde{\theta_6^9}|_{Y_7} = 0 - \left(-\dfrac{1}{y_9}\dfrac{\partial}{\partial x_9}\right) = \dfrac{\partial}{\partial y_{10}} \\ \delta(\theta_6)_{9\,11} = \tilde{\theta_6^{11}} - \tilde{\theta_6^9}|_{Y_7} = 0 - \left(-\dfrac{1}{y_9}\dfrac{\partial}{\partial x_9}\right) = \dfrac{1}{x_{11}}\dfrac{\partial}{\partial y_{11}} \end{cases}$$

Set $\{\tau_9 = 0, \tau_{10} = \frac{\partial}{\partial y_{10}}, \tau_{11} = 0\} \in C^0(N_{Y_7/S})$.

$$\begin{cases} \delta(\theta_6)_{9\,10} - \eta_7^{9\,10} = \dfrac{\partial}{\partial y_{10}} = \tau_{10} - \tau_9 \\ \delta(\theta_6)_{9\,11} - \eta_7^{9\,11} = 0 = \tau_{11} - \tau_9 \end{cases}$$

Hence we have $\{\delta(\theta_6)_{9\,10}, \delta(\theta_6)_{9\,11}\} = \eta_7$.

∘ $\delta(\theta_7) = \eta_6 - 2\eta_7 + \eta_8$

$$\delta(\theta_7)_{89} = \tilde{\theta_7^9} - \tilde{\theta_7^8}|_{Y_6} = \frac{1+ty_9}{x_9}\frac{\partial}{\partial y_9} - 0 = \frac{1}{x_9}\frac{\partial}{\partial y_9} = \eta_6^{89}$$

$$\begin{cases} \delta(\theta_7)_{9\,10} = \tilde{\theta_7^{10}} - \tilde{\theta_7^9}|_{Y_7} = -\dfrac{t+x_{10}}{y_{10}}\dfrac{\partial}{\partial x_{10}} - \dfrac{1+ty_9}{x_9}\dfrac{\partial}{\partial y_9} = -\dfrac{t+x_{10}}{x_{10}}\dfrac{\partial}{\partial y_{10}} \\ \delta(\theta_7)_{9\,11} = \tilde{\theta_7^{11}} - \tilde{\theta_7^9}|_{Y_7} = -\dfrac{1}{y_{11}}\dfrac{\partial}{\partial x_{11}} - \dfrac{1+ty_9}{x_9}\dfrac{\partial}{\partial y_9} = \dfrac{t-2x_{11}}{x_{11}(-t+x_{11})}\dfrac{\partial}{\partial y_{11}} \end{cases}$$

$$\begin{cases} \delta(\theta_7)_{11\,12} = \tilde{\theta_7^{12}} - \tilde{\theta_7^{11}}|_{Y_8} = 0 - \left(-\dfrac{1}{y_{11}}\dfrac{\partial}{\partial x_{11}}\right) = \dfrac{\partial}{\partial y_{12}} = \eta_{11\,12} \\ \delta(\theta_7)_{11\,13} = \tilde{\theta_7^{13}} - \tilde{\theta_7^{11}}|_{Y_8} = 0 - \left(-\dfrac{1}{y_{11}}\dfrac{\partial}{\partial x_{11}}\right) = \dfrac{1}{x_{13}}\dfrac{\partial}{\partial y_{13}} = \eta_8^{11\,13} \end{cases}$$

Set $\{\tau_9 = t\frac{\partial}{\partial x_9}, \tau_{10} = -\frac{\partial}{\partial y_{10}}, \tau_{11} = 0\} \in C^0(N_{Y_7/S})$.

$$\begin{cases} \delta(\theta_7)_{9\,10} + 2\eta_7^{9\,10} = -\dfrac{t+x_{10}}{x_{10}}\dfrac{\partial}{\partial y_{10}} = \tau_{10} - \tau_9 \\ \delta(\theta_7)_{9\,11} + 2\eta_7^{9\,11} = \dfrac{t}{x_{11}(t-x_{11})}\dfrac{\partial}{\partial y_{11}} = \tau_{11} - \tau_9 \end{cases}$$

Hence we have $\{\delta(\theta_7)_{9\,10}, \delta(\theta_7)_{9\,11}\} = -2\eta_7$.

$\circ$ $\delta(\theta_8) = \eta_7 - 2\eta_8$

$$\begin{cases} \delta(\theta_8)_{9\,10} = \tilde{\theta}_8^{10} - \tilde{\theta}_8^{9}|_{Y_7} = 0 - 0 = \eta_7^{9\,10} \\ \delta(\theta_8)_{9\,11} = \tilde{\theta}_8^{11} - \tilde{\theta}_8^{9}|_{Y_7} = \dfrac{1+a_0y_{11}}{x_{11}}\dfrac{\partial}{\partial y_{11}} - 0 = \eta_7^{9\,11} \end{cases}$$

$$\begin{cases} \delta(\theta_8)_{11\,12} = \tilde{\theta}_8^{12} - \tilde{\theta}_8^{11}|_{Y_8} = -\dfrac{a_0+x_{12}}{y_{12}}\dfrac{\partial}{\partial x_{12}} - \dfrac{1+a_0y_{11}}{x_{11}}\dfrac{\partial}{\partial y_{11}} = -\dfrac{a_0+x_{12}}{x_{12}}\dfrac{\partial}{\partial y_{12}} \\ \delta(\theta_8)_{11\,13} = \tilde{\theta}_8^{13} - \tilde{\theta}_8^{11}|_{Y_8} = -\dfrac{1}{y_{13}}\dfrac{\partial}{\partial x_{13}} - \dfrac{1+a_0y_{11}}{x_{11}}\dfrac{\partial}{\partial y_{11}} = \dfrac{-a_0+2x_{13}}{x_{13}(a_0-x_{13})}\dfrac{\partial}{\partial y_{13}} \end{cases}$$

Set $\{\tau_{11} = a_0\frac{\partial}{\partial x_{11}}, \tau_{12} = \frac{\partial}{\partial y_{12}}, \tau_{13} = 0\} \in C^0(N_{Y_8/S})$.

$$\begin{cases} \delta(\theta_8)_{11\,12} + 2\eta_{11\,12} = \dfrac{-a_0+x_{12}}{x_{12}}\dfrac{\partial}{\partial y_{12}} = \tau_{12} - \tau_{11} \\ \delta(\theta_8)_{11\,13} + 2\eta_8^{11\,13} = \dfrac{a_0}{x_{13}(a_0-x_{13})}\dfrac{\partial}{\partial y_{13}} = \tau_{13} - \tau_{11} \end{cases}$$

Hence we have $\{\delta(\theta_8)_{11\,12}, \delta(\theta_8)_{11\,13}\} = -2\eta_8$.

Summing up all the computations, we see that the matrix of the linear map δ is given by

$$\begin{pmatrix} -2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -2 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & -2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -2 \end{pmatrix}.$$

Since it is well-known that this matrix coincides with the intersection matrix $((Y_i, Y_j))_{1\le i,j\le r}$ of type $\tilde{E}_8$, this completes the proof of Proposition 2.1. $\square$

4. LOCAL COHOMOLOGY OF GENERALIZED OKAMOTO–PAINLEVÉ PAIR OF MULTIPLICATIVE TYPE

Let (S, Y) be a generalized Okamoto-Painlevé pair as in §2.

For a pair (S, Y) of additive type, the result of §2 shows the existence of differential equations on $S - Y$ (cf. [STT]). Even for (S, Y) of multiplicative type, if $\dim H^1_D(\Theta_S(-\log D)) \ge 1$, we can derive a differential equation in the same way as in [STT]. Unfortunately, we can prove $H^1_D(\Theta_S(-\log D)) = \{0\}$ for a pair (S, Y) of $\tilde{A}_8$. (For other multiplicative types, we expect that $H^1_D(\Theta_S(-\log D)) = \{0\}$.) This means that, for a pair (S, Y) of multiplicative type, there is no differential equation on $S - Y$ in the sense as in [STT].

In this section, we will calculate the local cohomology group $H^1_D(\Theta_S(-\log D))$ of pair (S, Y) of $\tilde{A}_8$ type.

- Construction of (S, Y) of $\tilde{A}_8$ type

Now, we consider (S, Y) of $\tilde{A}_8$ type as an example of multiplicative type.
According to [Sakai], any $\tilde{A}_8$-surface is obtained by blowing up $\mathbf{P}^2$ at the following 9 points given by


$z = 0$
$x = 0$
$p_7 \leftarrow p_8 \leftarrow p_9$
$p_1 \leftarrow p_2 \leftarrow p_6$
$p_5 \rightarrow p_4 \rightarrow p_3$
$y = 0$

$$p_1 : (1:0:0) \leftarrow p_2 : \left(\frac{z}{x}, \frac{y}{z}\right) = (0,0) \leftarrow p_6 : \left(\frac{z}{x}, \frac{xy}{z^2}\right) = (0,1),$$

$$p_3 : (0:0:1) \leftarrow p_4 : \left(\frac{y}{z}, \frac{x}{y}\right) = (0,0) \leftarrow p_5 : \left(\frac{y}{z}, \frac{zx}{y^2}\right) = (0,b),$$

$$p_7 : (0:1:0) \leftarrow p_8 : \left(\frac{x}{y}, \frac{z}{x}\right) = (0,0) \leftarrow p_9 : \left(\frac{x}{y}, \frac{yz}{x^2}\right) = (0,c).$$

Moreover there exists a $\mathbf{C}^\times$-action on the family of surfaces by

$$(\mu, (b, c, [x:y:z])) \mapsto (\mu^3 b, \mu^{-3} c, [\mu x : \mu^{-1} y : z]).$$

By putting $t = bc$, we can normalize this description.

We can choose the following coordinate system of $\tilde{A}_8$-surfaces S_t parameterized by t.



$E_i \simeq \mathbf{P}^1,\ E_i^2 = -1,$

$Y_i \simeq \mathbf{P}^1,\ Y_i^2 = -2.$

$$\begin{aligned}
U_i &= \operatorname{Spec}\mathbf{C}\,[x_i, y_i] \ \cong\ \mathbf{C}^2 \quad (i = 0, 1, ..., 8).\\
U_9 &= \operatorname{Spec}\mathbf{C}\left[x_9, y_9, \frac{1}{t + x_9}\right] \ \cong\ \mathbf{C}^2 - \{t + x_9 = 0\}.\\
U_i &= \operatorname{Spec}\mathbf{C}\left[x_i, y_i, \frac{1}{1 + x_i}\right] \ \cong\ \mathbf{C}^2 - \{1 + x_i = 0\} \quad (i = 10, 11).\\
U_{12} &= \operatorname{Spec}\mathbf{C}\left[u_{12}, v_{12}, \frac{1}{t + u_{12}v_{12}}\right] \ \cong\ \mathbf{C}^2 - \{t + u_{12}v_{12} = 0\}.\\
U_i &= \operatorname{Spec}\mathbf{C}\left[u_i, v_i, \frac{1}{1 + u_i v_i}\right] \ \cong\ \mathbf{C}^2 - \{1 + u_i v_i = 0\} \quad (i = 13, 14).
\end{aligned}$$

where $t \in \mathbf{C}^{\times}$.

$$\begin{aligned}
&Y_1 = \{x_0 = 0, y_1 = 0\}, && Y_2 = \{x_1 = 0, y_2 = 0, y_9 = 0\},\\
&Y_3 = \{x_2 = 0, y_3 = 0\}, && Y_4 = \{x_3 = 0, y_4 = 0\},\\
&Y_5 = \{x_4 = 0, y_5 = 0, y_{10} = 0\}, && Y_6 = \{x_5 = 0, y_6 = 0\},\\
&Y_7 = \{x_6 = 0, y_7 = 0\}, && Y_8 = \{x_7 = 0, y_8 = 0, y_{11} = 0\},\\
&Y_9 = \{x_8 = 0, y_0 = 0\}, &&
\end{aligned}$$

$$S_t = \bigcup_{i=0}^{14} U_i,$$

$$Y_t = \sum_{i=1}^{9} Y_i, \quad D = Y,$$

$$S_t - Y_t = U_{12} \cup U_{13} \cup U_{14}.$$

For later use, we need only the coordinate transformations near each component Y_i. Here we will list up the coordinate transformations only for a neighborhood of each Y_i.

$$Y_1 : \begin{cases} x_0 = {x_1}^2 y_1 \\ y_0 = \dfrac{1}{x_1} \end{cases} \qquad Y_2 : \begin{cases} x_1 = x_2 y_2 \\ y_1 = \dfrac{1}{x_2} \end{cases} \begin{cases} x_1 = x_9(t + x_9) y_9 \\ y_1 = \dfrac{1}{(t + x_9)} \end{cases} \qquad Y_3 : \begin{cases} x_2 = {x_3}^2 y_3 \\ y_2 = \dfrac{1}{x_3} \end{cases}$$

$$Y_4 : \begin{cases} x_3 = {x_4}^2 y_4 \\ y_3 = \dfrac{1}{x_4} \end{cases} \qquad Y_5 : \begin{cases} x_4 = x_5 y_5 \\ y_4 = \dfrac{1}{x_5} \end{cases} \begin{cases} x_4 = x_{10}(1 + x_{10}) y_{10} \\ y_4 = \dfrac{1}{(1 + x_{10})} \end{cases} \qquad Y_6 : \begin{cases} x_5 = {x_6}^2 y_6 \\ y_5 = \dfrac{1}{x_6} \end{cases}$$

$$Y_7 : \begin{cases} x_6 = {x_7}^2 y_7 \\ y_6 = \dfrac{1}{x_7} \end{cases} \qquad Y_8 : \begin{cases} x_7 = x_8 y_8 \\ y_7 = \dfrac{1}{x_8} \end{cases} \begin{cases} x_7 = x_{11}(1 + x_{11}) y_{11} \\ y_7 = \dfrac{1}{(1 + x_{11})} \end{cases} \qquad Y_9 : \begin{cases} x_8 = {x_0}^2 y_0 \\ y_8 = \dfrac{1}{x_0}. \end{cases}$$

This gives a generalized Okamoto–Painlevé pair (S_t, Y_t) of type $\tilde{A}_8$.

Proposition 4.1. *Let* (S_t, Y_t) *be as above and set* $D_t = (Y_t)_{red}$.
If $-t \in \mathbf{C}$ *is not a root of unity, then we have*

$$H^1_{D_t}(\Theta_{S_t}(-\log D_t)) = 0.$$

Remark 4.1. We expect that, if $-t$ is a root of unity then (S_t, Y_t) is of fibered type.

Proof. From the diagram

$$\begin{tikzcd}
 & & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & \Theta_S(-\log D) \arrow[r] \arrow[d, equal] & \Theta_S(-\log D) \otimes \mathcal{O}_S((n-1)D) \arrow[r] \arrow[d] & \Theta_S(-\log D) \otimes N_{(n-1)D} \arrow[r] \arrow[d, "\mu"] & 0 \\
0 \arrow[r] & \Theta_S(-\log D) \arrow[r] & \Theta_S(-\log D) \otimes \mathcal{O}_S(nD) \arrow[r] \arrow[d] & \Theta_S(-\log D) \otimes N_{nD} \arrow[r] \arrow[d] & 0 \\
 & & \Theta_S(-\log D) \otimes N_D^{\otimes n} \arrow[r, phantom, "\simeq"] \arrow[d] & \operatorname{coker} \mu \arrow[d] & \\
 & & 0 & 0 &
\end{tikzcd}$$

,

we obtain the exact sequence

$$0 \longrightarrow \Theta_S(-\log D) \otimes N_{(n-1)D} \longrightarrow \Theta_S(-\log D) \otimes N_{nD} \longrightarrow \Theta_S(-\log D) \otimes N_D^{\otimes n} \longrightarrow 0.$$

Therefore, we have the following sequence for each $n \geq 2$.

$$0 \longrightarrow H^0(\Theta_S(-\log D) \otimes N_{(n-1)D}) \longrightarrow H^0(\Theta_S(-\log D) \otimes N_{nD}) \longrightarrow H^0(\Theta_S(-\log D) \otimes N_D^{\otimes n}).$$

If $H^0(\Theta_S(-\log D) \otimes N_D^{\otimes n}) = 0$ for any $n \geq 1$, then we have

$$H^0(\Theta_S(-\log D) \otimes N_{(n-1)D}) \simeq H^0(\Theta_S(-\log D) \otimes N_{nD}).$$

Since $H^0(\Theta_S(-\log D) \otimes N_D) = 0$, we have

$$H^0(\Theta_S(-\log D) \otimes N_{nD}) = 0 \quad (n \geq 1).$$

Therefore, noting that (cf. §2)

$$H^1_D(S, \Theta_S(-\log D)) = \varinjlim H^0(\Theta_S(-\log D) \otimes N_{nD}),$$

we obtain

$$H^1_D(S, \Theta_S(-\log D)) = 0.$$

Now let us calculate $H^0(\Theta_S(-\log D) \otimes N_D^{\otimes n})$.

Recall that we have the exact sequence

$$0 \longrightarrow \nu_*(\oplus_{i=1}^9 N_{Y_i}) \longrightarrow \Theta_S(-\log D) \otimes N_D \longrightarrow \Theta_D \otimes N_D \longrightarrow 0,$$

where $\nu : \tilde{D} = \coprod_{i=1}^9 Y_i \to D$ is the normalization map (Lemma 2.1 (3)). Tensoring this sequence with $N_D^{\otimes n-1}$, we obtain

$$0 \longrightarrow \nu_*(\oplus_{i=1}^9 N_{Y_i}) \otimes N_D^{\otimes n-1} \longrightarrow \Theta_S(-\log D) \otimes N_D^{\otimes n} \longrightarrow \Theta_D \otimes N_D^{\otimes n} \longrightarrow 0.$$

Therefore we have

$$\begin{aligned}&H^0(D, \nu_*(\oplus_{i=1}^9 N_{Y_i}) \otimes N_D^{\otimes n-1}) \to H^0(D, \Theta_S(-\log D) \otimes N_D^{\otimes n}) \to H^0(D, \Theta_D \otimes N_D^{\otimes n}) \to \\ &H^1(D, \nu_*(\oplus_{i=1}^9 N_{Y_i}) \otimes N_D^{\otimes n-1}).\end{aligned}$$

In $\tilde{A}_8$ case, we see that $t_i = 2$, where t_i is the number of intersections of Y_i with other components. From this, we obtain

$$\Theta_D \simeq \nu_*(\oplus_{i=1}^9 \Theta_{Y_i}(-t_i)) \simeq \nu_*(\oplus_{i=1}^9 \mathcal{O}_{Y_i}(2-t_i)) \simeq \nu_*(\oplus_{i=1}^9 \mathcal{O}_{Y_i}).$$

Since ν is a finite morphism, we have

$$\begin{aligned}&H^i(D, \nu_*(\oplus_{i=1}^9 N_{Y_i}) \otimes N_D^{\otimes n-1}) \simeq H^i(\tilde{D}, (\oplus_{i=1}^9 N_{Y_i}) \otimes \nu^*(N_D^{\otimes n-1})) \quad (i = 0, 1), \\ &H^0(D, \Theta_D \otimes N_D^{\otimes n}) \simeq H^0(D, \nu_*(\oplus_{i=1}^9 \mathcal{O}_{Y_i}) \otimes N_D^{\otimes n}) \simeq H^0(\tilde{D}, (\oplus_{i=1}^9 \mathcal{O}_{Y_i}) \otimes \nu^*(N_D^{\otimes n})).\end{aligned}$$

Note that $D \cdot Y_i = 2 - t_i = 0$, we see that $N_D|_{Y_i} \simeq \mathcal{O}_{Y_i}$. Therefore we have

$$(\oplus_{i=1}^9 N_{Y_i}) \otimes \nu^*(N_D^{\otimes n-1}) \simeq \oplus_{i=1}^9 N_{Y_i}, \quad (\oplus_{i=1}^9 \mathcal{O}_{Y_i}) \otimes \nu^*(N_D^{\otimes n}) \simeq \oplus_{i=1}^9 \mathcal{O}_{Y_i}.$$

Summing up the arguments above, we obtain

$$\begin{tikzcd} 0 \arrow[r] & H^0(\Theta_S(-\log D)\otimes N_D^{\otimes n}) \arrow[r] & \oplus_{i=1}^9 H^0(Y_i,\mathcal{O}_{Y_i}) \arrow[r, "\delta"] \arrow[d, equal] & \oplus_{i=1}^9 H^1(N_{Y_i}) \arrow[d, equal] \\ & & \mathbf{C}^9 \arrow[r, "\delta"] & \mathbf{C}^9. \end{tikzcd}$$

We will analyse the edge homomorphism δ by using the Čech cocycles.

Noting that

$$H^0(\Theta_D\otimes N_D^{\otimes n})\simeq H^0(\Theta_{Y_i}(-2)\otimes N_D^{\otimes n})\simeq \mathbf{C}^9. \tag{6}$$

For each i $(1\le i\le 9)$, we introduce a generator θ_i of the cohomology group in (6) as follows.

θ_1	$\left\{\theta_1^0=\frac{y_0}{x_0^n y_0^n}\frac{\partial}{\partial y_0}\text{ on }U_0\cap Y_1,\quad \theta_1^1=-\frac{x_1}{x_1^n y_1^n}\frac{\partial}{\partial x_1}\text{ on }U_1\cap Y_1\right\}$
θ_2	$\left\{\begin{array}{c}\theta_2^1=\frac{(1-ty_1)^n y_1}{x_1^n y_1^n}\frac{\partial}{\partial y_1}\text{ on }U_1\cap Y_2,\quad \theta_2^2=-\frac{(x_2-t)^n x_2}{x_2^n y_2^n}\frac{\partial}{\partial x_2}\text{ on }U_2\cap Y_2,\\ \theta_2^9=-\frac{1}{(t+x_9)^{n-1}y_9^n}\frac{\partial}{\partial x_9}\text{ on }U_9\cap Y_2\end{array}\right\}$
θ_3	$\left\{\theta_3^2=\frac{y_2}{x_2^n y_2^n}\frac{\partial}{\partial y_2}\text{ on }U_2\cap Y_3,\quad \theta_3^3=-\frac{x_3}{x_3^n y_3^n}\frac{\partial}{\partial x_3}\text{ on }U_3\cap Y_3\right\}$
θ_4	$\left\{\theta_4^3=\frac{y_3}{x_3^n y_3^n}\frac{\partial}{\partial y_3}\text{ on }U_3\cap Y_4,\quad \theta_4^4=-\frac{x_4}{x_4^n y_4^n}\frac{\partial}{\partial x_4}\text{ on }U_4\cap Y_4\right\}$
θ_5	$\left\{\begin{array}{c}\theta_5^4=\frac{(1-x_4)^n y_4}{x_4^n y_4^n}\frac{\partial}{\partial y_4}\text{ on }U_4\cap Y_5,\quad \theta_5^5=-\frac{(x_5-1)^n x_5}{x_5^n y_5^n}\frac{\partial}{\partial x_5}\text{ on }U_5\cap Y_5,\\ \theta_5^{10}=-\frac{1}{(1+x_{10})^{n-1}y_{10}^n}\frac{\partial}{\partial x_{10}}\text{ on }U_{10}\cap Y_5\end{array}\right\}$
θ_6	$\left\{\theta_6^5=(-1)^n\frac{y_5}{x_5^n y_5^n}\frac{\partial}{\partial y_5}\text{ on }U_5\cap Y_6,\quad \theta_6^6=-(-1)^n\frac{x_6}{x_6^n y_6^n}\frac{\partial}{\partial x_6}\text{ on }U_6\cap Y_6\right\}$
θ_7	$\left\{\theta_7^6=(-1)^n\frac{y_6}{x_6^n y_6^n}\frac{\partial}{\partial y_6}\text{ on }U_6\cap Y_7,\quad \theta_7^7=-(-1)^n\frac{x_7}{x_7^n y_7^n}\frac{\partial}{\partial x_7}\text{ on }U_7\cap Y_7\right\}$
θ_8	$\left\{\begin{array}{c}\theta_8^7=(-1)^n\frac{(1-x_7)^n y_7}{x_7^n y_7^n}\frac{\partial}{\partial y_7}\text{ on }U_7\cap Y_8,\quad \theta_8^8=-(-1)^n\frac{(x_8-1)^n x_8}{x_8^n y_8^n}\frac{\partial}{\partial x_8}\text{ on }U_8\cap Y_8,\\ \theta_8^{11}=-(-1)^n\frac{1}{(1+x_{11})^{n-1}y_{11}^n}\frac{\partial}{\partial x_{11}}\text{ on }U_{11}\cap Y_8\end{array}\right\}$
θ_9	$\left\{\theta_9^8=\frac{y_8}{x_8^n y_8^n}\frac{\partial}{\partial y_8}\text{ on }U_8\cap Y_9,\quad \theta_9^0=-\frac{x_0}{x_0^n y_0^n}\frac{\partial}{\partial x_0}\text{ on }U_0\cap Y_9\right\}$

On the other hand, for each i $(1\le i\le 9)$, we have a generator $\eta_i\in H^1(Y_i,N_{Y_i/S}\otimes N_D^{\otimes n-1})$ as follows.

η_1	$\left\{\eta_1^{01} = \frac{1}{x_1 x_1^{n-1} y_1^{n-1}} \frac{\partial}{\partial y_1} \text{ on } U_0 \cap\ U_1 \cap Y_1\right\}$
η_2	$\left\{\eta_2^{12} = \frac{1}{y_2^{n-1}} \frac{\partial}{\partial y_2} \text{ on } U_1 \cap\ U_2 \cap Y_2,\ \eta_2^{29} = 0 \text{ on } U_2 \cap\ U_9 \cap Y_2\right\}$
η_3	$\left\{\eta_3^{23} = \frac{1}{x_3 x_3^{n-1} y_3^{n-1}} \frac{\partial}{\partial y_3} \text{ on } U_2 \cap\ U_3 \cap Y_3\right\}$
η_4	$\left\{\eta_4^{34} = \frac{1}{x_4 x_4^{n-1} y_4^{n-1}} \frac{\partial}{\partial y_4} \text{ on } U_3 \cap\ U_4 \cap Y_4\right\}$
η_5	$\left\{\eta_5^{45} = \frac{1}{y_5^{n-1}} \frac{\partial}{\partial y_5} \text{ on } U_4 \cap\ U_5 \cap Y_5,\ \eta_5^{5\,10} = 0 \text{ on } U_5 \cap\ U_{10} \cap Y_5\right\}$
η_6	$\left\{\eta_6^{56} = (-1)^n \frac{1}{x_6 x_6^{n-1} y_6^{n-1}} \frac{\partial}{\partial y_6} \text{ on } U_5 \cap\ U_6 \cap Y_6\right\}$
η_7	$\left\{\eta_7^{67} = (-1)^n \frac{1}{x_7 x_7^{n-1} y_7^{n-1}} \frac{\partial}{\partial y_7} \text{ on } U_6 \cap\ U_7 \cap Y_7\right\}$
η_8	$\left\{\eta_8^{78} = (-1)^n \frac{1}{y_8^{n-1}} \frac{\partial}{\partial y_8} \text{ on } U_7 \cap\ U_8 \cap Y_8,\ \eta_8^{8\,11} = 0 \text{ on } U_8 \cap\ U_{11} \cap Y_8\right\}$
η_9	$\left\{\eta_9^{80} = \frac{1}{x_0 x_0^{n-1} y_0^{n-1}} \frac{\partial}{\partial y_0} \text{ on } U_8 \cap\ U_0 \cap Y_9\right\}$

We take $\{\theta_i\}$ and $\{\eta_i\}$ as basis of $H^0(\Theta_D \otimes N_D^{\otimes n})$ and $\oplus_{i=1}^{9} H^1(N_{Y_i/S} \otimes N_D^{\otimes n-1})$ respectively. By using these bases, we compute the matrix representing the connecting homomorphisn δ.

For that purpose, let us lift 0-cocycle θ_1 to 0-cochains of $\Theta_S(-\log D) \otimes N_D$ as

$$\tilde{\theta_1^0} = \frac{y_0}{x_0^n y_0^n} \frac{\partial}{\partial y_0} \text{ on } U_0, \quad \tilde{\theta_1^1} = -\frac{x_1}{x_1^n y_1^n} \frac{\partial}{\partial x_1} \text{ on } U_1,$$

$$\tilde{\theta_1^i} = 0 \text{ on } U_i \ (i = 2, 3, \cdots, 14).$$

Other 0-cocycles can be lifted in a similar way.

We first compute $\delta(\theta_1)$.

From the definition of δ, we have $\delta(\theta_1) = \{\delta(\theta_1)_{ij} \text{ on } U_i \cap U_j \cap D\}$ with

$$\delta(\theta_1)_{80} = -\tilde{\theta_1^8} + \tilde{\theta_1^0}|_{Y_9} = -0 + \frac{y_0}{x_0^n y_0^n} \frac{\partial}{\partial y_0} = \eta_9^{80}$$

$$\delta(\theta_1)_{01} = -\tilde{\theta_1^0} + \tilde{\theta_1^1}|_{Y_1} = -\frac{y_0}{x_0^n y_0^n} \frac{\partial}{\partial y_0} + \left(-\frac{x_1}{x_1^n y_1^n} \frac{\partial}{\partial x_1}\right) = -2 \frac{1}{x_1 x_1^{n-1} y_1^{n-1}} \frac{\partial}{\partial y_1} = -2\eta_1^{01}$$

$$\begin{cases} \delta(\theta_1)_{12} = -\tilde{\theta_1^1} + \tilde{\theta_1^2}|_{Y_2} = -\left(-\frac{x_1}{x_1^n y_1^n} \frac{\partial}{\partial x_1}\right) + 0 = \frac{1}{y_2^{n-1}} \frac{\partial}{\partial y_2} = \eta_2^{12} \\ \delta(\theta_1)_{29} = -\tilde{\theta_1^2} + \tilde{\theta_1^9}|_{Y_2} = -0 + 0 = \eta_2^{29} \end{cases}$$

Other $\delta(\theta_1)_{ij}$'s are zero.

Obviously $\delta(\theta_1) = \eta_9 - 2\eta_1 + \eta_2$.

Other $\delta(\theta_i)$'s can be treated in a similar way. In what follows, we just list up a few results of computations.

∘ $\delta(\theta_2) = \eta_1 - 2\eta_2 + (-t)^n\eta_3$

$$\delta(\theta_2)_{01} = -\tilde{\theta}_2^0 + \tilde{\theta}_2^1|_{Y_1} = -0 + \frac{(1-ty_1)^n y_1}{x_1^n y_1^n}\frac{\partial}{\partial y_1} = \frac{1}{x_1 x_1^{n-1} y_1^{n-1}}\frac{\partial}{\partial y_1} = \eta_1^{01} \ (y_1 = 0 \text{ on } U_1 \cap Y_1)$$

$$\begin{cases} \delta(\theta_2)_{12} = -\tilde{\theta}_2^1 + \tilde{\theta}_2^2|_{Y_2} = -\dfrac{(1-ty_1)^n y_1}{x_1^n y_1^n}\dfrac{\partial}{\partial y_1} + \left(-\dfrac{(x_2-t)^n x_2}{x_2^n y_2^n}\dfrac{\partial}{\partial x_2}\right) = -\dfrac{(x_2-t)^n}{x_2 x_2^{n-1} y_2^{n-1}}\dfrac{\partial}{\partial y_2} \\ \delta(\theta_2)_{29} = -\tilde{\theta}_2^2 + \tilde{\theta}_2^9|_{Y_2} = -\left(-\dfrac{(x_2-t)^n x_2}{x_2^n y_2^n}\dfrac{\partial}{\partial x_2}\right) + \left(-\dfrac{1}{(t+x_9)^{n-1} y_9^n}\dfrac{\partial}{\partial x_9}\right) = -\dfrac{1}{x_9 (t+x_9)^{n-1} y_9^{n-1}}\dfrac{\partial}{\partial y_9} \end{cases}$$

$$\delta(\theta_2)_{23} = -\tilde{\theta}_2^2 + \tilde{\theta}_2^3|_{Y_3} = -\left(-\frac{(x_2-t)^n x_2}{x_2^n y_2^n}\frac{\partial}{\partial x_2}\right)\frac{\partial}{\partial x_2} + 0 = (-t)^n \frac{1}{x_3 x_3^{n-1} y_3^{n-1}}\frac{\partial}{\partial y_3} = (-t)^n \eta_3^{23}$$

Set $\{\tau_1 = \frac{(1-ty_1)^n + (1-ty_1)^{n-1} - 2}{y_1 x_1^{n-1} y_1^{n-1}}\frac{\partial}{\partial x_1}, \tau_2 = \frac{(x_2-t)^{n-1}}{x_2^{n-1} y_2^{n-1}}\frac{\partial}{\partial y_2}, \tau_9 = 0\} \in C^0(N_{Y_2/M} \otimes N_D^{\otimes n-1})$.

$$\begin{cases} \delta(\theta_2)_{12} + 2\eta_2^{12} = -\dfrac{(x_2-t)^n}{x_2 x_2^{n-1} y_2^{n-1}}\dfrac{\partial}{\partial y_2} + 2\dfrac{1}{y_2^{n-1}}\dfrac{\partial}{\partial y_2} = -\tau_1 + \tau_2 \\ \delta(\theta_2)_{29} + 2\eta_2^{29} = -\dfrac{1}{x_9 (t+x_9)^{n-1} y_9^{n-1}}\dfrac{\partial}{\partial y_9} - 0 = -\tau_2 + \tau_9 \end{cases}$$

Thus we have $\{\delta(\theta_2)_{12}, \delta(\theta_2)_{29}\} = -2\eta_2$.

∘ $\delta(\theta_3) = \frac{1}{(-t)^n}\eta_2 - 2\eta_3 + \eta_4$

$$\begin{cases} \delta(\theta_3)_{12} = -\tilde{\theta}_3^1 + \tilde{\theta}_3^2|_{Y_2} = -0 + \dfrac{y_2}{x_2^n y_2^n}\dfrac{\partial}{\partial y_2} = \dfrac{y_2}{x_2^n y_2^n}\dfrac{\partial}{\partial y_2} \\ \delta(\theta_3)_{29} = -\tilde{\theta}_3^2 + \tilde{\theta}_3^9|_{Y_2} = -\dfrac{y_2}{x_2^n y_2^n}\dfrac{\partial}{\partial y_2} + 0 = -\dfrac{1}{x_9^n (t+x_9)^n y_9^{n-1}}\dfrac{\partial}{\partial y_9} \end{cases}$$

$$\delta(\theta_3)_{23} = -\tilde{\theta}_3^2 + \tilde{\theta}_3^3|_{Y_3} = -\frac{y_2}{x_2^n y_2^n}\frac{\partial}{\partial y_2} + \left(-\frac{x_3}{x_3^n y_3^n}\frac{\partial}{\partial x_3}\right) = -2\frac{1}{x_3 x_3^{n-1} y_3^{n-1}}\frac{\partial}{\partial y_3} = -2\eta_3^{23}$$

$$\delta(\theta_3)_{34} = -\tilde{\theta}_3^3 + \tilde{\theta}_3^4|_{Y_4} = -\left(-\frac{x_3}{x_3^n y_3^n}\frac{\partial}{\partial x_3}\right) + 0 = \frac{1}{x_4 x_4^{n-1} y_4^{n-1}}\frac{\partial}{\partial y_4} = \eta_4^{34}$$

Set $\{\tau_1 = \frac{1-(1-ty_1)^n}{(-t)^n y_1 x_1^{n-1} y_1^{n-1}}\frac{\partial}{\partial x_1}, \tau_2 = \frac{(-t)^n - (x_2-t)^n}{(-t)^n x_2 x_2^{n-1} y_2^{n-1}}\frac{\partial}{\partial y_2}, \tau_9 = -\frac{1}{(-t)^n (x_9+t)^n y_9^{n-1}}\frac{\partial}{\partial y_9}\} \in C^0(N_{Y_2/M} \otimes N_D^{\otimes n-1})$.

$$\begin{cases} \delta(\theta_3)_{12} - \dfrac{1}{(-t)^n}\eta_2^{12} = \dfrac{y_2}{x_2^n y_2^n}\dfrac{\partial}{\partial y_2} - \dfrac{1}{(-t)^n}\dfrac{1}{y_2^{n-1}}\dfrac{\partial}{\partial y_2} = -\tau_1 + \tau_2 \\ \delta(\theta_3)_{29} - \dfrac{1}{(-t)^n}\eta_2^{29} = -\dfrac{1}{x_9^n (t+x_9)^n y_9^{n-1}}\dfrac{\partial}{\partial y_9} - 0 = -\tau_2 + \tau_9 \end{cases}$$

Thus we have $\{\delta(\theta_3)_{12}, \delta(\theta_3)_{29}\} = \frac{1}{(-t)^n}\eta_2$.

Summing up all the computations, we see that the matrix of the linear map δ is given by

$$\delta = \begin{pmatrix} -2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & -2 & (-t)^{-n} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & (-t)^n & -2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -2 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -2 \end{pmatrix}$$

Since

$$\det \delta = \frac{((-t)^n - 1)^2}{(-t)^n},$$

we see

$$\operatorname{rank} \delta = \begin{cases} 9 & (-t \text{ is not a root of unity}) \\ 8 & (-t \text{ is a root of unity}). \end{cases}$$

Therefore, if $-t$ is not a root of unity, then

$$H^0(\Theta_S(-\log D) \otimes N_D^{\otimes n}) = 0.$$

□

Acknowledgements

The author is deeply grateful to Professor Masa-Hiko Saito, who gave support to the whole of this work. We would also like to thank Kenji Iohara for many useful advices.

References

[B-W] D. M. Burns, Jr. & J. M. Wahl, Local contributions to global deformations of surfaces, Invent. Math. **26** (1974), 67-88.

[Gr] A. Grothendieck, *Local cohomology*, (noted by R. Hartshorne), Lecture Notes in Math. 41, Springer-Verlag, Berlin, Heidelberg, New York (1967), 106 pp.

[Kaw] Y. Kawamata, On deformations of compactifiable manifolds, Math. Ann., **235**, (1978), 247–265.

[O1] K. Okamoto, Sur les feuilletages associés sux équ ations du second ordre à points critiques fixes de P. Painlevé, Es paces des conditions initiales, Japan. J. Math., **5**, 1979, 1–79.

[O2] K. Okamoto, Polynomial Hamiltonians associated with Painlevé equations, I, II, Proc. Japan Acad., **56**, (1980), 264–268; ibid, 367–371.

[O3] K. Okamoto, Studies on the Painlevé equations I. Annali di Mathematica pura ed applicata **CXLVI** 1987, 337–381; II. Japan. J. Math., **13**, (1987), 47–76; III. Math. Ann. **275** (1986), 221–255; IV. Funkcial. Ekvac. Ser. Int. **30** (1987), 305- -332.

[Sa-Tak] M.-H. Saito & T. Takebe, Classification of Okamoto–Painlevé pairs, preprint, Kobe 2000.

[STT] M.-H. Saito , T. Takebe & H. Terajima, Deformation of Okamoto–Painlevé pair and Painlevé equations, preprint, Kobe 2000.

[Sakai] H. Sakai, Rational surfaces associated with affine root systems and geometry of the Painlevé equations, preprint, Kyoto-Math 99-10.

Department of Mathematics, Faculty of Science, Kobe University, Kobe, Rokko, 657-8501, Japan
E-mail address: `terajima@math.kobe-u.ac.jp`